

## METHODS OF APPLICATION



Apply as  
Irrigation  
or Fertigation



Manual Application



Irrigation Systems



Watering Systems

## APPLICATION GUIDE

### Specific Rates

| Crop Type  | Dilution in Water | Application Rates  | Application Growth Stage   |
|--|-------------------|--|--|
| <b>Leafy vegetables</b> (e.g. Broccoli, Cabbage, Cauliflower, Celery, Herbs, Lettuce, Kohlrabi, Rhubarb, Spinach, Spring Onion)  | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>5L/ha</b>  | Early Growth<br>Mid-season<br>Onset of Maturity                        |
| <b>Fruiting vegetables</b> (e.g. Bean, Capsicum, Corn, Cucumber, Eggplant, Melons, Pumpkin, Sweet pea, Squash, Tomato, Zucchini) | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>5L/ha</b>  | Early vegetative growth<br>Mid vegetative growth<br>Early fruit growth |
| <b>Bulbs and root crops</b> (e.g. Beet, Carrot, Garlic, Onion, Parsnip, Potato, Radish, Turnip, Sweet potato)                    | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>5L/ha</b>  | Early vegetative growth<br>Mid vegetative growth<br>Early Bulking      |
| <b>Fruit trees</b> (e.g. Apple, Apricot, Cherry, Citrus, Fig, Nectarines, Olives, Peaches, Pear, Persimmon, Plums)               | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>20L/ha</b> | 2-4 weeks before budburst<br>Pre-flowering<br>Soon after harvest       |
| <b>Grapes, Soft fruits</b> (Strawberries and Berries)  | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>5L/ha</b>  | 2-4 weeks before budburst<br>Pre-flowering<br>Soon after harvest       |
| <b>Young trees or Non-bearing fruit tree trees</b>   | x 100             | 1st application at <b>20L/ha</b><br>2nd application at <b>20L/ha</b><br>3rd application at <b>20L/ha</b> | 2-4 weeks before budburst<br>Mid season<br>Late season                 |

### Application Guide

Fertigation products should be applied at the end of the irrigation cycle to prevent fertiliser front going below the root zone. After injection, the system should run long enough to clear the fertiliser out of the lines.

## HOW TO MIX



Shake  
Vigorously



Mix  
with Water



Mix with  
other Chemicals



### PRODUCT COMPATIBILITY + JAR TESTING

DO NOT mix with alkaline copper fungicides or inoculants. If you are unsure, we recommend a simple jar test of products. Mix together and check if reaction occurs.



### PRECAUTIONS

Non-toxic product. Avoid unneeded contact. Keep out of the reach of children. If contact is made with eyes, immediately rinse with plenty of water. If swallowed, seek medical attention.

## ANALYSIS AND PRODUCT ASSURANCE

RLF



Australian-owned Formulator, Manufacturer and Supplier of High-analysis Broad-spectrum Liquid Fertiliser technologies. For over 25 years RLF's products have been used by millions of farmers and growers world-wide. ISO 9001 Quality Assured Company since 1998.



### MACRO NUTRIENTS

Nitrogen (N)  
Phosphorus (P)  
Phosphorus ( $P_2O_5$ )  
Potassium (K)  
Potassium ( $K_2O$ )  
Sulphur (S)  
Magnesium (Mg)

### MICRO NUTRIENTS

Zinc (Zn)  
Manganese (Mn)  
Copper (Cu)  
Iron (Fe)  
Boron (B)  
Molybdenum (Mo)

### GLOBAL

%w/w  
%w/w  
%w/w  
%w/w  
%w/w  
%w/w

%w/w  
%w/w  
%w/w  
%w/w  
%w/w  
%w/w



### Member Login

Please login to be able to view this detail



Not a member yet?  
[Register Here](#)

[LOG IN](#)