


METHODS OF APPLICATION


To the Leaf



Foliar Spray onto the Crop Leaf



Manual Application




Machine Application



Rain Safe in 2 hours

To the Soil



Furrow Injection

APPLICATION GUIDE

Specific Rates

Dilution in Water		Application Rates	Recommended Timings
Minimum	Maximum		
1 to 30	1 to 100	1st application at 2L/ha at about 30% banding	4-6 leaf stage
1 to 30	1 to 100	2nd application at 4L/ha broadcast or desirable banding	at Early squaring
1 to 30	1 to 100	3rd application at 5L/ha	at Early flowering


Furrow Injection Rate and Dilution

Dilution in Water		Application Rates
Minimum	Maximum	
x40	- x100	2 - 3 Litres/hectare (L/ha)

HOW TO MIX



Shake Vigorously



Mix with Water



Mix with other Chemicals



PRODUCT COMPATIBILITY + JAR TESTING

DO NOT mix with alkaline copper fungicides or inoculants. If unsure about compatibility, we recommend a simple jar test of products. Mix together and check if reaction occurs.



PRECAUTIONS

Non-toxic product. Avoid unneeded contact. Keep out of the reach of children. If contact is made with eyes, immediately rinse with plenty of water. If swallowed, seek medical attention.

ANALYSIS AND PRODUCT ASSURANCE

RLF

Australian-owned Formulator, Manufacturer and Supplier of High-analysis Broad-spectrum Liquid Fertiliser technologies. For over 25 years RLF's products have been used by millions of farmers and growers world-wide. ISO 9001 Quality Assured Company since 1998.



MACRO NUTRIENTS

Nitrogen (N)
Phosphorus (P)
Phosphorus (P)
Potassium (K)
Potassium (K)
Sulphur (S)
Magnesium (Mg)

MICRO NUTRIENTS

Zinc (Zn)
Manganese (Mn)
Copper (Cu)
Iron (Fe)
Boron (B)
Cobalt (Co)
Molybdenum (Mo)

GLOBAL

%w/w
%w/w
%w/w
%w/w
%w/w
%w/w
%w/w

%w/w
%w/w
%w/w
%w/w
%w/w
%w/w
%w/w



Member Login

Please login to be able to view this detail

Not a member yet?
[Register Here](#)

[LOG IN](#)