

Foliar Fertiliser

METHODS OF APPLICATION



Foliar Fertiliser to Spray onto the Crop Leaf



Manual Application



Machine Application



Rain Safe in 2 hours

APPLICATION GUIDE

Specific Rates

Crop Type	Carrier Rate Litres/hectare	Application Rate (Litres/hectare)		
Wheat (all Cereals) Corn Canola Dryland Pasture Hay Fodder Crops (oats, millet, sorghum, turnip and other forage brassicas) Fruit trees and Vegetables	1L to 10 L to 1L to 20L	4 - 6 Litres/hectare (L/ha)	Note: 2-3 weeks is required before foliar application can be repeated	

Recommended Timings

Neconimental rinings						
Сгор Туре	Number of Minimum	Applications Preferred	Timing for Application			
Wheat (all Cereals)	1 to	2 times	Good canopy formation into grain filling (avoid anthesis)			
Corn	1 to	2 times	Good canopy formation into grain filling (avoid silking)	10		
Canola	1 to	2 times	Good ground cover to early flowering	10		
Dryland Pasture	1 to	2 times	Good ground cover after each grazing in winter or early spring			
Hav	1 to	2 times	Good ground cover when shut for hay or silage			
Fodder Crops (oats, millet, sorghum, turnip and other forage brassicas)	1 to	2 times	Good ground cover and after each grazing when re-growth is expected			
Fruit trees and Vegetables	1 to	2 times	Good ground cover and after each grazing when re-growth is expected			



Fertigation via Irrigation or **Sprinkler Systems**



Manual Application



Irrigation Systems



Crop Type	Litres / ha per Irrigation	Number of Applications per season / year	
Young Vnes, Olives and Citrus trees Mature Vines Mature Olives & Citrus trees Other mature Fruit Trees Vegetable Crops Irrigated Pastures	10 Litres 20 Litres 20 Litres 20 Litres 20 Litres 20-30 Litres	Bimonthly to monthly Bud burst and before flowering Before flowering and post harvest Up to flowering and after harvest Early vegetative growth and as required After each cut or grazing or as required	

ном то міх



Shake Vigorously



Mix with Water



Mix with other Chemicals



PRODUCT COMPATIBILITY + JAR TESTING

DO NOT mix with alkaline copper fungicides or inoculants. If you are unsure, we recommend a simple jar test of products. Mix together and check if reaction occurs.

PRECAUTIONS

Non-toxic product. Avoid unneeded contact. Keep out of the reach of children. If contact is made with eyes, immediately rinse with plenty of water. If swallowed, seek medical attention.

ANALYSIS AND PRODUCT ASSURANCE

RLF

Australian-owned Formulator, Manufacturer and Supplier of High-analysis Broad-spectrum Liquid Fertiliser technologies. For over 25 years RLF's products have been used by millions of farmers and growers world-wide. ISO 9001 Quality Assured Company since 1998.



MACRO NUTRIENT

Potassium (as K₂O)

MICRO NUTRIENT

Silica (as SiO₂)



Please login to be able to view this detail

Not a member yet? Register Here

LOG IN

