

Fertigation Fertiliser

METHODS OF APPLICATION



Apply as Irrigation or Fertigation







Irrigation Systems



APPLICATION GUIDE

Specific Rates

| Сгор Туре | Minimum Dilution in Water | Application Rates | Application Growth Stage |
|--|---------------------------|---------------------------|---------------------------|
| eafy vegetables (e.g. Broccoli, Cabbage, | x 100 | 1st application at 20L/ha | Early Growth |
| Cauliflower, Celery, Herbs, Lettuce, Kohlrabi, | | 2nd application at 20L/ha | Mid-season |
| Rhubarb, Spinach, Spring Onion) | | 3rd application at 5L/ha | Onset of Maturity |
| ruiting vegetables (e.g. Bean, Capsicum, | x 100 | 1st application at 20L/ha | Early vegetative growth |
| Corn, Cucumber, Eggplant, Melons, Pumpkin, | | 2nd application at 20L/ha | Mid vegetative growth |
| Sweet pea, Squash, Tomato, Zucchini) | | 3rd application at 5L/ha | During fruit growth |
| Bulbs and root crops (e.g. Beet, Carrot, | x 100 | 1st application at 20L/ha | Early vegetative growth |
| Garlic, Onion, Parsnip, Potato, Radish, Turnip, | 5/20/04/05/44 | 2nd application at 20L/ha | Mid vegetative growth |
| weet potato) | | 3rd application at 5L/ha | Early Bulking |
| ruit trees (e.g. Apple, Apricot, Cherry, Citrus, | x 100 | 1st application at 20L/ha | Soon after harvest |
| Fig, Nectarines, Olives, Peaches, Pear, | Coesses | 2nd application at 20L/ha | 2-4 weeks before budburst |
| ersimmon, Plums) | | 3rd application at 5L/ha | During fruit growth |
| Grapes, Soft fruits (Strawberries and Berries) | x 100 | 1st application at 20L/ha | 2-4 weeks before budburst |
| | | 2nd application at 20L/ha | Pre-flowering |
| | | 3rd application at 5L/ha | During berry growth |
| | | 4th application at 20L/ha | Soon after harvest |
| Young trees or Non-bearing fruit tree trees | x 100 | 1st application at 20L/ha | 2-4 weeks before budburst |
| | | 2nd application at 20L/ha | Mid season |
| | | 3rd application at 20L/ha | Late season |

Application Guide

Fertigation products should be applied at the end of the irrigation cycle to prevent fertiliser front going below the root zone. After injection, the system should run long enough to clear the fertiliser out of the lines.

HOW TO MIX



Shake Vigorously



Mix with Water



other Chemicals



Mix with



PRODUCT COMPATIBILITY + JAR TESTING

DO NOT mix with alkaline copper fungicides or inoculants. If you are unsure, we recommend a simple jar test of products. Mix together and check if reaction occurs.



PRECAUTIONS

Non-toxic product. Avoid unneeded contact. Keep out of the reach of children. If contact is made with eyes, immediately rinse with plenty of water. If swallowed, seek medical attention.

ANALYSIS AND PRODUCT ASSURANCE

RLF

Australian-owned Formulator, Manufacturer and Supplier of High-analysis Broad-spectrum Liquid Fertiliser technologies. For over 25 years RLF's products have been used by millions of farmers and growers world-wide. ISO 9001 Quality Assured Company since 1998.



MACRO NUTRIENTS Nitrogen (N)

Phosphorus (P) Phosphorus (POS) Potassium (K) Potassium (K2O)



Member Login

Please login to be able to view this detail

Not a member yet? Register Here

LOG IN

BAL

w/w

w/w

w/w

w/w

W/W

