

METHODS OF APPLICATION




**Foliar Fertiliser to
Spray onto the
Crop Leaf**



Manual Application



Machine Application



Rain Safe in 2 hours

APPLICATION GUIDE

Specific Rates

| Crop Type | Carrier Rate Litres/hectare/L of product | Application Rate (Litres/hectare) |
|---|--|-----------------------------------|
| Wheat (all Cereals) | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |
| Corn | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |
| Canola | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |
| Dryland Pasture | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |
| Hay | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |
| Fodder Crops (oats, millet, sorghum, turnip and other forage brassicas) | 10 - 20 X | 4 - 6 Litres/hectare (L/ha) |

Note :

2-3 weeks is required before foliar application can be repeated

Recommended Timings


| Crop Type | Number of Applications Minimum Preferred | Timing for Application |
|---|---|---|
| Wheat (all Cereals) | 1 to 2 times | Good canopy formation into grain filling (avoid anthesis) |
| Corn | 1 to 2 times | Good canopy formation into grain filling (avoid silking) |
| Canola | 1 to 2 times | Good ground cover to early flowering |
| Dryland Pasture | 1 to 2 times | Good ground cover after each grazing in winter or early spring |
| Hay | 1 to 2 times | Good ground cover when shut for hay or silage |
| Fodder Crops (oats, millet, sorghum, turnip and other forage brassicas) | 1 to 2 times | Good ground cover and after each grazing when re-growth is expected |



**Fertigation
via Irrigation or
Sprinkler Systems**



Manual Application




Irrigation Systems




Watering Systems

| Crop Type | Litres / ha per Irrigation | Number of Applications per season / year |
|--------------------------------------|----------------------------|--|
| Young Vines, Olives and Citrus trees | 10 Litres | Bimonthly to monthly |
| Mature Vines | 20 Litres | Bud burst and before flowering |
| Mature Olives & Citrus trees | 20 Litres | Before flowering and post harvest |
| Other mature Fruit Trees | 20 Litres | Up to flowering and after harvest |
| Vegetable Crops | 20 Litres | Early vegetative growth and as required |
| Irrigated Pastures | 20-30 Litres | After each cut or grazing or as required |

HOW TO MIX



**Shake
Vigorously**



**Mix
with Water**



**Mix with
other Chemicals**



PRODUCT COMPATIBILITY + JAR TESTING

DO NOT mix with alkaline copper fungicides or inoculants. If you are unsure, we recommend a simple jar test of products. Mix together and check if reaction occurs.



PRECAUTIONS

Non-toxic product. Avoid unneeded contact. Keep out of the reach of children. If contact is made with eyes, immediately rinse with plenty of water. If swallowed, seek medical attention.

ANALYSIS AND PRODUCT ASSURANCE

RLF



Australian-owned Formulator, Manufacturer and Supplier of High-analysis Broad-spectrum Liquid Fertiliser technologies. For over 25 years RLF's products have been used by millions of farmers and growers world-wide. ISO 9001 Quality Assured Company since 1998.



TYPICAL ANALYSIS

Calcium (Ca)
Nitrogen (N)
Magnesium (Mg)
Zinc (Zn)
Manganese (Mn)
Copper (Cu)
Iron (Fe)
Boron (B)
Molybdenum (Mo)



Member Login

Please login to be able to view this detail

Not a member yet?
[Register Here](#)

[LOG IN](#)